

RESEARCH

Open Access

Bounds on the number of closed walks in a graph and its applications

Xiaodan Chen¹ and Jianguo Qian^{2*}

*Correspondence:

jqqian@xmu.edu.cn

²School of Mathematical Sciences,
Xiamen University, Siming Road
422, Xiamen, 361005, P.R. China
Full list of author information is
available at the end of the article

Abstract

Using graph-theoretical techniques, we establish an inequality regarding the number of walks and closed walks in a graph. This inequality yields several upper bounds for the number of closed walks in a graph in terms of the number of vertices, number of edges, maximum degree, degree sequence, and the Zagreb indices of the graph. As applications, we also present some new upper bounds on the Estrada index for general graphs, bipartite graphs, trees and planar graphs, some of which improve the known results obtained by using the algebraic techniques.

MSC: 05C50; 94C15; 05C38

Keywords: number of closed walks; Estrada index; Zagreb indices; maximum degree; degree sequence

1 Introduction

Throughout this paper we consider simple graphs, *i.e.*, graphs without loops and multiple edges. For a graph G with n ($n \geq 2$) vertices, the adjacency matrix of G is, as usual, defined as the $n \times n$ matrix $A(G) = [a_{ij}]$ in which $a_{ij} = 1$ if the i th and the j th vertices are adjacent, and $a_{ij} = 0$ otherwise. The eigenvalues of $A(G)$ are also called the eigenvalues of the graph G . For a vertex v in G , we denote by $N(v)$ and $d(v)$ the neighbor (the set of vertices adjacent to v) and the degree of v , respectively. The degree sequence of G , denoted by (d_1, d_2, \dots, d_n) , is a list of the vertex degrees of G in non-increasing order. Let $V(G)$ and $E(G)$ denote the vertex set and edge set of G , respectively. The first and the second Zagreb indices of G are defined as

$$Zg_1(G) = \sum_{v \in V(G)} d^2(v) \quad \text{and} \quad Zg_2(G) = \sum_{uv \in E(G)} d_u d_v,$$

respectively [1].

A walk W of length k starting at a vertex v_0 and ending at a vertex v_k in G is a sequence of vertices, *i.e.*, $v_0 v_1 v_2 \dots v_k$, in which v_i is adjacent to v_{i+1} for each $i = 0, 1, \dots, k-1$. In particular, if the vertices $v_0, v_1, v_2, \dots, v_k$ (except the possible v_0 and v_k) are pairwise distinct, then W is well known as a path, and if $v_0 = v_k$ then W is called a closed walk. It is well known [2] that the number of closed walks of length k in G is exactly the trace of $A(G)^k$ which, in turn, is the sum of the k th power of the eigenvalues of G (known as the k th spectral moment of G). This fact is of importance in the theory of total π -electron energy, for details see [3, 4] and the references cited therein. Also, the sequence of the numbers of

closed walks of length k , $k = 1, 2, \dots$, starting at a given vertex v , was proposed by Randić [5] for characterization of the environment of vertex v .

Based on the number of closed walks, Estrada [6] put forward a graph invariant, which was originally referred to as the subgraph centrality but has since become known as the Estrada index of a graph G , defined as

$$EE(G) = \sum_{i=1}^n e^{\lambda_i},$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of G . The Estrada index has successfully found applications in various fields, including biochemistry [6, 7] and complex networks [8]. Also, a number of mathematical properties, especially various lower and upper bounds on the Estrada index of a graph have been established, for details we refer the reader to [9–16]; other properties can be found in [17–19] and a latest survey paper by Gutman *et al.* [20].

In general, counting the closed walks in a graph (of large order) is not an easy work. Only a few results were obtained for some special types of graphs, *e.g.*, vertex-transitive graphs [21] and generalized de Bruijn graphs [22]. In this paper, using graph-theoretical techniques, we establish an inequality regarding the number of walks and closed walks starting at a given vertex. This inequality yields several upper bounds for the number of closed walks in a graph in terms of the number of vertices, number of edges, maximum degree, degree sequence, the first and the second Zagreb indices of the graph. As applications, in Section 3 we present some new upper bounds on the Estrada index for general graphs, bipartite graphs, trees, and planar graphs, which improve some known results obtained by using the algebraic techniques.

2 Main results

Given a graph G and a vertex v , let $\mathcal{W}_k(G, v)$ denote the set of walks of length k starting at v in G , and let $W_k(G, v) = |\mathcal{W}_k(G, v)|$. Obviously, $W_0(G, v) = 1$, $W_1(G, v) = d(v)$. Moreover, it is easy to check that

$$W_2(G, v) = \sum_{u \in N(v)} d(u),$$

$$W_3(G, v) = \sum_{u \in N(v)} \sum_{w \in N(u)} d(w).$$

In general, we have the following result.

Lemma 1 *Let G be a graph of order n with maximum degree Δ , and let v be an arbitrary vertex in G . Then*

$$W_k(G, v) \leq \Delta^k, \quad \text{for any } k \geq 1, \tag{1}$$

$$W_k(G, v) \leq d(v)\Delta^{k-1}, \quad \text{for any } k \geq 2, \tag{2}$$

$$W_k(G, v) \leq \sum_{u \in N(v)} d(u)\Delta^{k-2}, \quad \text{for any } k \geq 3, \tag{3}$$

$$W_k(G, v) \leq \sum_{u \in N(v)} \sum_{w \in N(u)} d(w) \Delta^{k-3}, \quad \text{for any } k \geq 4. \tag{4}$$

Each of the equalities holds in (1)-(4) for all v if and only if G is regular.

Proof Let $W = v_1 v_2 \cdots v_k$ be a walk in $\mathcal{W}_k(G, v)$. Observe that each of k steps of W has at most Δ choices, then (1) follows. We also notice that the first one, two, and three step(s) of W have exactly $d(v)$, $\sum_{u \in N(v)} d(u)$ and $\sum_{u \in N(v)} \sum_{w \in N(u)} d(w)$ choices, respectively, and each of the remaining steps has at most Δ choices, so (2), (3), and (4) follow as well. Moreover, it is not difficult to see that each of the equalities holds in (1)-(4) for all v in G if and only if G is a Δ -regular graph. This completes the proof. \square

Let $W_k(G)$ be the number of walks of length k in G , i.e., $W_k(G) = \sum_{v \in V(G)} W_k(G, v)$. It is clear that $W_0(G) = n$, $W_1(G) = \sum_{v \in V(G)} d(v) = 2m$. Moreover, one can deduce easily that

$$\begin{aligned} W_2(G) &= \sum_{v \in V(G)} \sum_{u \in N(v)} d(u) = \sum_{u \in V(G)} d^2(u) = Zg_1(G), \\ W_3(G) &= \sum_{v \in V(G)} \sum_{u \in N(v)} \sum_{w \in N(u)} d(w) = \sum_{u \in V(G)} d(u) \sum_{w \in N(u)} d(w) \\ &= 2 \sum_{uw \in E(G)} d(u)d(w) = 2Zg_2(G), \end{aligned}$$

where $Zg_1(G)$, $Zg_2(G)$ are the first and the second Zagreb indices of G , respectively. Using these facts and Lemma 1, for general $k \geq 1$, we have the following.

Theorem 2 *Let G be a graph with n vertices, m edges, and maximum degree Δ . Then*

$$W_k(G) \leq n \Delta^k, \quad \text{for any } k \geq 1, \tag{5}$$

$$W_k(G) \leq 2m \Delta^{k-1}, \quad \text{for any } k \geq 2, \tag{6}$$

$$W_k(G) \leq Zg_1(G) \Delta^{k-2}, \quad \text{for any } k \geq 3, \tag{7}$$

$$W_k(G) \leq 2Zg_2(G) \Delta^{k-3}, \quad \text{for any } k \geq 4. \tag{8}$$

Each of the equalities holds in (5)-(8) if and only if G is regular.

Proof This proof is trivial. \square

Remark Bounds (6)-(8) can be seen as some slight improvements of bound (5). Here, we also list two other improvements of bound (5) for using later.

(i) [23] Let G be a graph of order n with degree sequence (d_1, d_2, \dots, d_n) . Then for $k \geq 1$,

$$W_k(G) \leq \sum_{i=1}^n d_i^k, \tag{9}$$

with equality if and only if G is regular or $k \leq 2$.

- (ii) [24] Let G be a graph of order n with maximum degree Δ . If G admits an orientation with maximum outdegree $d \leq \Delta/2$, then for $k \geq 1$,

$$W_k(G) \leq n2^k d^{k/2} (\Delta - d)^{k/2}. \tag{10}$$

Moreover, from the proof of (10) (see Theorem 16 in [24]), one can deduce that the equality holds in (10) if and only if G is a Δ -regular Euler graph.

Now we turn to the number of closed walks. Let $CW_k(G, v)$ denote the set of closed walks of length k starting and ending at v in G , and let $CW_k(G, v) = |CW_k(G, v)|$. It is obvious that $0 = CW_1(G, v) < W_0(G, v) = 1$, $CW_2(G, v) = W_1(G, v) = d(v)$. In general, for $k \geq 3$ we establish the following simple but useful result.

Lemma 3 *Let G be a graph of order n and let v be an arbitrary vertex in G . Then, for any $k \geq 3$,*

$$CW_k(G, v) \leq W_{k-1}(G, v), \tag{11}$$

with equality if and only if k is even, and the component of G containing v is bipartite and v is adjacent to each of the vertices in the other partition part.

Proof Let f be the map from $CW_k(G, v)$ into $W_{k-1}(G, v)$ such that, for any closed walk $v\nu_1\nu_2 \cdots \nu_{k-1}v \in CW_k(G, v)$,

$$f(v\nu_1\nu_2 \cdots \nu_{k-1}v) = v\nu_1\nu_2 \cdots \nu_{k-1}.$$

One can see that different closed walks are mapped to different walks by f , which yields

$$CW_k(G, v) \leq W_{k-1}(G, v).$$

Further, the equality holds if and only if the end vertex of each walk in $W_{k-1}(G, v)$ is adjacent to v . In this case, if k is odd then, for any edge νu , $W_{k-1}(G, v)$ contains a walk of the form $W = vuvu \cdots vuv$ while v is not adjacent to itself, a contradiction. So if the equality holds in (11), then k must be even.

Now we consider the component C_v of G containing v , under the assumption that the equality holds in (11) and k is even. Let $N_l(G, v)$ denote the set of vertices at distance l from v in G . We claim that $N_l(G, v)$ is an empty set, for any $l \geq 3$. Otherwise there would be a path $P = v\nu_1\nu_2\nu_3$ such that ν_3 is not adjacent to v , but the walk $W = v\nu_1\nu_1 \cdots v\nu_1\nu_2\nu_3$ defined on the path P belongs to $W_{k-1}(G, v)$, which implies that ν_3 is adjacent to v , a contradiction. We next show that there are no edges with both end vertices in $N_l(G, v)$, $l \in \{1, 2\}$. For contradiction, assume that there is an edge, say $\nu_1\nu_2$, with $\nu_1, \nu_2 \in N_1(G, v)$. Then $T = v\nu_1\nu_2\nu_1$ is a triangle and consequently the walk $W = v\nu_1\nu_1 \cdots v\nu_1\nu_2\nu_1$ defined on T belongs to $W_{k-1}(G, v)$, which implies that v is adjacent to v , again a contradiction. Similarly, if there are some edges with both end vertices in $N_2(G, v)$, then there must exist a path $P = v\nu_1\nu_2\nu_3$ such that ν_3 is not adjacent to v , which also yields a contradiction as the above argument. Thus, it follows that the component C_v is bipartite with partition $(N_1(G, v), N_2(G, v) \cup \{v\})$.

The converse is obvious, completing the proof. \square

Let $CW_k(G)$ denote the number of closed walks of length k in G , i.e., $CW_k(G) = \sum_{v \in V(G)} CW_k(G, v)$. Clearly, $CW_0(G) = n$, $CW_1(G) = 0$ and $CW_2(G) = 2m$. For any $k \geq 3$, using Lemma 3, we have

Theorem 4 *Let G be a graph of order n with maximum degree Δ and degree sequence (d_1, d_2, \dots, d_n) . Then, for any $k \geq 3$,*

$$CW_k(G) \leq n\Delta^{k-1}, \tag{12}$$

$$CW_k(G) \leq 2m\Delta^{k-2}, \tag{13}$$

$$CW_k(G) \leq Zg_1(G)\Delta^{k-3}, \tag{14}$$

$$CW_k(G) \leq 2Zg_2(G)\Delta^{k-4} \quad (\text{for } k \geq 4), \tag{15}$$

$$CW_k(G) \leq \sum_{i=1}^n d_i^{k-1}. \tag{16}$$

Each of the equalities holds in (12)-(16) if and only if k is even and each component of G is the complete bipartite graph $K_{\Delta, \Delta}$. Moreover, if G admits an orientation with maximum outdegree $d \leq \Delta/2$, then, for any $k \geq 3$,

$$CW_k(G) \leq n2^{k-1}d^{(k-1)/2}(\Delta - d)^{(k-1)/2}, \tag{17}$$

with equality if and only if both k and Δ are even and each component of G is the complete bipartite graph $K_{\Delta, \Delta}$.

Proof For $k \geq 3$, it follows from Lemma 3 that

$$CW_k(G) = \sum_{v \in V(G)} CW_k(G, v) \leq \sum_{v \in V(G)} W_{k-1}(G, v) = W_{k-1}(G),$$

with equality if and only if k is even, and each component of G is a complete bipartite graph. This result together with bounds (5)-(10) yield bounds (12)-(17) directly; also the equality cases follow by noting that G is Δ -regular (Δ is even in the case of (17)). The proof is completed. \square

Recall that an orientation of a graph G is a digraph D obtained from G by choosing an orientation for each edge. The outdegree of a vertex v in D is the number of edges with tail v . It is well known [24] that a tree (or forest) admits an orientation with maximum outdegree $d = 1$ and a planar graph with $d = 3$. In fact, for a forest, fixing a root for each component and orienting each edge in each component toward its root would yield an orientation with $d = 1$; furthermore, a planar graph has an orientation with $d = 3$ since its edges can be partitioned into three forests (see, e.g., [25]). Thus, by (17) we get an immediate corollary.

Corollary 5 *Let G be a graph of order n with maximum degree Δ . If G is a tree (or forest) and $\Delta \geq 2$, then, for any $k \geq 3$,*

$$CW_k(G) < n(\sqrt{4(\Delta - 1)})^{k-1}. \tag{18}$$

If G is a planar graph and $\Delta \geq 6$, then, for any $k \geq 3$,

$$CW_k(G) < n(\sqrt{12(\Delta - 3)})^{k-1}. \tag{19}$$

Remark that if G is a bipartite graph (including tree and forest), then there are no closed walks of odd length in G , and hence $CW_k(G) = 0$ when k is odd. Formally this is stated in the following proposition.

Proposition 6 *Let G be a bipartite graph. Then, for any $k \geq 1$, $CW_{2k-1}(G) = 0$.*

3 Applications

In this section we apply the results in the previous section to estimate the Estrada index of graphs.

Let $M_k(G)$ denote the k th spectral moment of G , i.e., $M_k(G) = \sum_{i=1}^n \lambda_i^k$, where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of G . Then $M_k(G) = CW_k(G)$ [2]. On the other hand, recalling the power-series expansion of the function e^x , we have another expression for the Estrada index of G as follows:

$$EE(G) = \sum_{k=0}^{\infty} \frac{M_k(G)}{k!} = \sum_{k=0}^{\infty} \frac{CW_k(G)}{k!}. \tag{20}$$

In particular, if G is a bipartite graph, then by Proposition 6, we get

$$EE(G) = \sum_{k=0}^{\infty} \frac{M_{2k}(G)}{(2k)!} = \sum_{k=0}^{\infty} \frac{CW_{2k}(G)}{(2k)!}. \tag{21}$$

We are now ready to give some new upper bounds for $EE(G)$.

Theorem 7 *Let G be a graph with n vertices, m edges, t triangles, maximum degree Δ and degree sequence (d_1, d_2, \dots, d_n) . Then*

$$EE(G) < \frac{n}{\Delta} (e^\Delta - 1), \tag{22}$$

$$EE(G) < n + \frac{2m}{\Delta^2} (e^\Delta - 1 - \Delta), \tag{23}$$

$$EE(G) < n + m + \frac{Zg_1(G)}{\Delta^3} \left(e^\Delta - 1 - \Delta - \frac{\Delta^2}{2} \right), \tag{24}$$

$$EE(G) < n + m + t + \frac{2Zg_2(G)}{\Delta^4} \left(e^\Delta - 1 - \Delta - \frac{\Delta^2}{2} - \frac{\Delta^3}{6} \right), \tag{25}$$

$$EE(G) < \sum_{i=1}^n \frac{e^{d_i} - 1}{d_i}. \tag{26}$$

Moreover, if G admits an orientation with maximum outdegree $d \leq \Delta/2$, then

$$EE(G) < \frac{n}{\sqrt{4d(\Delta - d)}} (e^{\sqrt{4d(\Delta - d)}} - 1). \tag{27}$$

Proof We first consider (22). By (20), Theorem 4 and noticing that $CW_0(G) = n$, $CW_1(G) = 0$, $CW_2(G) = 2m \leq n\Delta$, we have

$$\begin{aligned} EE(G) &= CW_0(G) + CW_1(G) + \frac{CW_2(G)}{2!} + \sum_{k=3}^{\infty} \frac{CW_k(G)}{k!} \\ &< n + \frac{n\Delta}{2!} + \sum_{k=3}^{\infty} \frac{n\Delta^{k-1}}{k!} \\ &= n + \frac{n}{\Delta} \left(\sum_{k=0}^{\infty} \frac{\Delta^k}{k!} - 1 - \Delta \right) \\ &= \frac{n}{\Delta} (e^{\Delta} - 1). \end{aligned}$$

The discussion for (23)-(27) is analogous by observing that $CW_2(G) = 2m = \sum_{i=1}^n d_i$, $CW_3(G) = 6t < W_2(G) = Zg_1(G)$ and, $CW_2(G) = W_1(G) \leq n\sqrt{4d(\Delta - d)}$ with equality if and only if G is a Δ -regular Euler graph. \square

For bipartite graphs, from (21) and the power-series expansion of the hyperbolic cosine $\cosh(x) = (e^x + e^{-x})/2$, one can easily obtain the following result by a similar reasoning as in the proof of Theorem 7.

Theorem 8 *Let G be a bipartite graph with n vertices, m edges, maximum degree Δ and degree sequence (d_1, d_2, \dots, d_n) . Then*

$$EE(G) \leq n + \frac{n}{\Delta} (\cosh(\Delta) - 1), \tag{28}$$

$$EE(G) \leq n + \frac{2m}{\Delta^2} (\cosh(\Delta) - 1), \tag{29}$$

$$EE(G) \leq n + m + \frac{Zg_1(G)}{\Delta^3} \left(\cosh(\Delta) - 1 - \frac{\Delta^2}{2} \right), \tag{30}$$

$$EE(G) \leq n + m + \frac{2Zg_2(G)}{\Delta^4} \left(\cosh(\Delta) - 1 - \frac{\Delta^2}{2} \right), \tag{31}$$

$$EE(G) \leq n + \sum_{i=1}^n \frac{\cosh(d_i) - 1}{d_i}. \tag{32}$$

Each of the equalities holds in (28)-(32) if and only if each component of G is the complete bipartite graph $K_{\Delta, \Delta}$. Moreover, if G admits an orientation with maximum outdegree $d \leq \Delta/2$, then

$$EE(G) \leq n + \frac{n}{\sqrt{4d(\Delta - d)}} (\cosh(\sqrt{4d(\Delta - d)}) - 1), \tag{33}$$

with equality if and only if Δ is even and each component of G is the complete bipartite graph $K_{\Delta, \Delta}$.

Similar to Corollary 5, substituting $d = 1$ and $d = 3$ in (33) and (27), respectively, we have the following corollary.

Corollary 9 *Let G be a graph of order n with maximum degree Δ . If G is a tree (or forest) and $\Delta \geq 2$, then*

$$EE(G) < n + \frac{n}{\sqrt{4(\Delta-1)}} (\cosh(\sqrt{4(\Delta-1)}) - 1). \quad (34)$$

If G is a planar graph and $\Delta \geq 6$, then

$$EE(G) < \frac{n}{\sqrt{12(\Delta-3)}} (e^{\sqrt{12(\Delta-3)}} - 1). \quad (35)$$

Remark In the past few years, a number of upper bounds on the Estrada index of graphs have been established by using the algebraic techniques (see, for example, [10, 11, 14, 16]). In comparison to the algebraic techniques, the bounds based on our graph-theoretical method are related to the degree parameters (mainly the maximum degree), which are somewhat different from the previous ones. Moreover, our method would be more effective in some cases. For example, in [10] de la Peña *et al.* showed that, for any graph G ,

$$EE(G) \leq n - 1 + e^{\sqrt{2m}},$$

which was later improved by Zhou in [16] as follows:

$$EE(G) \leq n - 1 - \sqrt{2m} + e^{\sqrt{2m}}. \quad (36)$$

Here, our bound (23) would be better than the bound (36) when $2 \leq \Delta < \sqrt{2m}$. To see this, we first consider the function $f(x) = (e^x - 1 - x)/x^2$ with $x \geq 2$, which is strictly increasing with respect to x since $f'(x) = [(x-2)e^x + x + 2]/x^3 > 0$ when $x \geq 2$. Then for $2 \leq \Delta < \sqrt{2m}$, we get

$$n + \frac{2m}{\Delta^2} (e^\Delta - 1 - \Delta) = n + 2mf(\Delta) < n + 2mf(\sqrt{2m}) = n - 1 - \sqrt{2m} + e^{\sqrt{2m}}.$$

In addition, for a bipartite graph G , the authors in [10] proved that

$$EE(G) \leq n - 2 + 2 \cosh(\sqrt{m}). \quad (37)$$

Here, our bound (29) would be better than the bound (37) when $2 \leq \Delta < \sqrt{m}$. Indeed, when $x \geq 2$ the function $f(x) = (\cosh x - 1)/x^2$ is strictly increasing with respect to x since $f'(x) = [(x-2)(e^x - e^{-x}) + 4(1 - e^{-x})]/2x^3 > 0$. Therefore, for $2 \leq \Delta < \sqrt{m}$, we have

$$n + \frac{2m}{\Delta^2} (\cosh(\Delta) - 1) = n + 2mf(\Delta) < n + 2mf(\sqrt{m}) = n - 2 + 2 \cosh(\sqrt{m}).$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Author details

¹College of Mathematics and Information Science, Guangxi University, Daxue Road 100, Nanning, 530004, P.R. China.

²School of Mathematical Sciences, Xiamen University, Siming Road 422, Xiamen, 361005, P.R. China.

Acknowledgements

The authors would like to thank the anonymous referees for their extremely helpful comments and suggestions towards improving the original version of this paper. The first author was supported partially by the Scientific Research Foundation of Guangxi University (Grant No. XBZ130083) and NNSF of China (No. 11361007). The second author was supported by NNSF of China (No. 10831001).

Received: 29 September 2013 Accepted: 6 May 2014 Published: 20 May 2014

References

1. Gutman, I, Trinajstić, N: Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons. *Chem. Phys. Lett.* **17**, 535-538 (1972)
2. Cvetković, D, Doob, M, Sachs, H: *Spectra of Graphs-Theory and Application*. Academic Press, New York (1980)
3. Gutman, I: Remark on the moment expansion of total π -electron energy. *Theor. Chim. Acta* **83**, 313318 (1992)
4. Gutman, I, Marković, S, Vesović, A, Estrada, E: Approximating total π -electron energy in terms of spectral moments. A quantitative approach. *J. Serb. Chem. Soc.* **63**, 639-646 (1998)
5. Randić, M: Random walks and their diagnostic value for characterization of atomic environment. *J. Comput. Chem.* **1**, 386-399 (1980)
6. Estrada, E: Characterization of 3D molecular structure. *Chem. Phys. Lett.* **319**, 713-718 (2000)
7. Estrada, E, Rodríguez-Valázquez, JA, Randić, M: Atomic branching in molecules. *Int. J. Quant. Chem.* **106**, 823-832 (2006)
8. Estrada, E, Rodríguez-Valázquez, JA: Subgraph centrality in complex networks. *Phys. Rev. E* **71**, 056103 (2005)
9. Das, KC, Lee, SG: On the Estrada index conjecture. *Linear Algebra Appl.* **431**, 1351-1359 (2009)
10. de la Peña, JA, Gutman, I, Rada, J: Estimating the Estrada index. *Linear Algebra Appl.* **427**, 70-76 (2007)
11. Fath-Tabar, GH, Ashrafi, AR: New upper bounds for Estrada index of bipartite graphs. *Linear Algebra Appl.* **435**, 2607-2611 (2011)
12. Gutman, I, Estrada, E, Rodríguez-Valázquez, JA: On a graph-spectrum-based structure descriptor. *Croat. Chem. Acta* **80**, 151-154 (2007)
13. Gutman, I: Lower bounds for Estrada index. *Publ. Inst. Math. (Belgr.)* **83**, 1-7 (2008)
14. Liu, J, Liu, B: Bounds of the Estrada index of graphs. *Appl. Math. J. Chin. Univ. Ser. B* **25**, 325-330 (2010)
15. Zhao, H, Jia, Y: On the Estrada index of bipartite graph. *MATCH Commun. Math. Comput. Chem.* **61**, 495-501 (2009)
16. Zhou, B: On Estrada index. *MATCH Commun. Math. Comput. Chem.* **60**, 485-492 (2008)
17. Du, Z, Zhou, B: The Estrada index of unicyclic graphs. *Linear Algebra Appl.* **436**, 3149-3159 (2012)
18. Ilic, A, Stevanović, D: The Estrada index of chemical trees. *J. Math. Chem.* **47**, 305-314 (2010)
19. Zhang, J, Zhou, B, Li, J: On Estrada index of trees. *Linear Algebra Appl.* **434**, 215-223 (2011)
20. Gutman, I, Deng, H, Radenković, S: The Estrada index: an updated survey. In: Cvetković, D, Gutman, I (eds.) *Selected Topics on Applications of Graph Spectra*, pp. 155-174. *Math. Inst., Beograd* (2011)
21. Jajcaya, R, Malniã, A, Marušć, D: On the number of closed walks in vertex-transitive graphs. *Discrete Math.* **307**, 484-493 (2007)
22. Shibata, Y, Shirahata, M, Osawa, S: Counting closed walks in generalized de Bruijn graphs. *Inf. Process. Lett.* **49**, 135-138 (1994)
23. Fiol, MA, Garriga, E: Number of walks and degree powers in a graph. *Discrete Math.* **309**, 2613-2614 (2009)
24. Hayes, TP: A simple condition implying rapid mixing of single-site dynamics on spin systems. In: 47th Ann. IEEE Symp. Found. Comp. Sci. (FOCS'06), pp. 39-46 (2006)
25. Gonçalves, D: Covering planar graphs with forests, one having bounded maximum degree. *J. Comb. Theory, Ser. B* **99**, 314-322 (2009)

10.1186/1029-242X-2014-199

Cite this article as: Chen and Qian: Bounds on the number of closed walks in a graph and its applications. *Journal of Inequalities and Applications* 2014, **2014**:199

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com